

In the Claims

For the convenience of the Examiner, Applicants submit herewith a complete claim set.
No new amendments to the claims have been made.

1-37. (Cancelled)

38. (Previously Presented) A method of identifying chemical substances which have potential pharmacological activity using nematode worms, which method comprises the steps of:

(a) dispensing substantially equal numbers of nematode worms into each of the wells of a multi-well assay plate;

(b) contacting the nematode worms with a chemical substance;

(c) detecting a signal indicating phenotypic, physiological, behavioral, or biochemical changes in the nematode worms using non-visual detection means;

wherein step (a) is performed in a multi-well plate with liquid assay medium containing a water soluble polymer at a concentration sufficient to increase the viscosity of the medium;

wherein the water soluble polymer is medium viscosity carboxymethyl cellulose.

39. (Previously Presented) The method as claimed in claim 38 wherein the concentration of water soluble polymer in the liquid medium is 0.3% (weight/volume).

40. (Previously Presented) A method of identifying chemical substances which have potential pharmacological activity using nematode worms, which method comprises the steps of:

(a) dispensing substantially equal numbers of nematode worms into each of the wells of a multi-well assay plate;

(b) contacting the nematode worms with a chemical substance;

(c) detecting a signal indicating phenotypic, physiological, behavioral, or biochemical changes in the nematode worms using non-visual detection means;

wherein step (a) is performed in a multi-well plate with liquid assay medium containing a water soluble polymer at a concentration sufficient to prevent the nematode worms from sticking to the wells of the multi-well plate.

41. (Previously Presented) The method as claimed in claim 40 wherein the water soluble polymer is polyethylene glycol, polyvinyl alcohol, or polyvinylpyrrolidone.

42. (Previously Presented) A method as claimed in claim 40 wherein the concentration of water soluble polymer in the liquid medium is from 0.01% to 10% (weight/volume).

43. (Previously Presented) A method as claimed in claim 42 wherein the concentration of water soluble polymer in liquid medium is 0.1% (weight/volume).

44-186. (Cancelled)

187. (Original) A method of identifying chemical substances which have potential pharmacological activity using nematode worms, which method comprises the steps of:

- (a) dispensing substantially equal numbers of nematode worms into each of the wells of a multi-well assay plate;
- (b) contacting the nematode worms with a chemical substance;
- (c) detecting a signal indicating phenotypic, physiological, behavioural, or biochemical changes in the nematode worms using non-visual detection means, wherein the assay is carried out in a liquid medium having a viscosity greater than M9 medium.

188. (Original) The method as claimed in claim 187 wherein the nematode worms are microscopic nematodes.

189. (Original) The method as claimed in claim 188 wherein the nematode worms are *C. elegans* or *C. briggsae*.

190. (Original) The method as claimed in claim 187 wherein the step of detecting a signal comprises detecting a change in a measurable property of a marker molecule, whereby a change

in the property of the marker molecule indicates a phenotypic, physiological, behavioural, or biochemical change in the nematode worms.

191. (Original) The method as claimed in claim 190 wherein the marker molecule is a fluorescent molecule, a luminescent molecule, or a coloured molecule.

192. (Original) The method as claimed in claim 190 wherein the marker molecule is a precursor of a fluorescent molecule, a precursor of a luminescent molecule, or a precursor of a coloured molecule.

193. (Original) The method as claimed in claim 192 wherein said marker molecule is capable of being cleaved by the action of an enzyme present in the gut of *C. elegans* to generate a fluorescent molecule, a luminescent molecule, or a coloured molecule.

194. (Original) The method as claimed in claim 187 wherein the non-visual detection means is a multi-well plate reader.

195. (Original) The method as claimed in claim 194 wherein the multi-well plate reader performs luminescence, fluorescence, or spectrophotometric detection.

196. (Original) The method as claimed in claim 187 wherein the non-visual detection means is a fluorescence activated nematode screening and sorting (FANS) device.

197. (Original) The method as claimed in claim 196 wherein the FANS device performs luminescence, fluorescence, or spectrophotometric detection.

198. (Original) The method as claimed in claim 187 wherein step (a) comprises dispensing substantially equal volumes of a homogeneous suspension of nematode worms into each of the wells of the multi-well assay plate.

199. (Original) The method as claimed in claim 198 wherein the homogeneous suspension comprises a suspension of *C. elegans* in a viscous solution.
200. (Original) The method as claimed in claim 199 wherein the viscous solution comprises a solution of a polymer material.
201. (Original) The method as claimed in claim 200 wherein the polymer material is low melting point agarose.
202. (Original) The method as claimed in claim 187 wherein the nematode worms are synchronized in the same growth stage.
203. (Original) The method as claimed in claim 202 wherein the nematode worms are eggs, L1 stage, L2 stage, L3 stage, L4 stage, adult worms, or dauer worms.
204. (Original) The method as claimed in claim 202 wherein the worms are hermaphrodites or males.
205. (Original) The method as claimed in claim 187 wherein the nematode worms are a wild type strain, a mutant strain, a transgenic strain, or a humanized strain.
206. (Original) The method as claimed in claim 187 wherein step (a) is performed in a multi-well plate with liquid assay medium containing a water soluble polymer at a concentration sufficient to increase the viscosity of the medium.
207. (Original) The method as claimed in claim 206 wherein the water soluble polymer is carboxymethyl cellulose, low melting point agarose, or polyethylene glycol.
208. (Original) The method as claimed in claim 207 wherein the water soluble polymer is medium viscosity carboxymethyl cellulose.

209. (Previously Presented) The method as claimed in claim 206 wherein the concentration of water soluble polymer in the liquid medium is 0.3% (weight/volume).

210. (Original) The method as claimed in claim 187 wherein step (a) is performed in a multi-well plate with liquid assay medium containing a water soluble polymer at a concentration sufficient to prevent the nematode worms from sticking to the wells of the multi-well plate.

211. (Previously Presented) The method as claimed in claim 210 wherein the water soluble polymer is polyethylene glycol, polyvinyl alcohol, or polyvinylpyrrolidone.

212. (Previously Presented) The method as claimed in claim 210 wherein the concentration of water soluble polymer in the liquid medium is from 0.01% to 10% (weight/volume).

213. (Previously Presented) The method as claimed in claim 212 wherein the concentration of water soluble polymer in the liquid medium is 0.1% (weight/volume).